

REFERENCER til artikel I Dansk Sportsmedicin nr. 4, 2014:

Styrketræning i rehabilitering

Jens Bojsen-Møller¹ og Per Aagaard²

¹Seksjon for fysisk prestasjonsevne, Norges Idrettshøgskole, Oslo, Norge

²Institut for Idræt og Biomekanik, Syddansk Universitet, Odense, Danmark

1. Suetta C, Aagaard P, Rosted A, Jakobsen AK, Duus B, Kjaer M, Magnusson SP. Training-induced changes in muscle CSA, muscle strength, EMG, and rate of force development in elderly subjects after long-term unilateral disuse. *Journal of Applied Physiology*. 2004;97(5):1954-61.
2. Aagaard P. Training-induced changes in neural function. *Exercise and Sport Sciences Reviews*. 2003;31(2):61-7.
3. Aagaard P, Suetta C, Caserotti P, Magnusson SP, Kjaer M. Role of the nervous system in sarcopenia and muscle atrophy with aging: strength training as a countermeasure. *Scandinavian Journal of Medicine & Science in Sports*. 2010;20(1):49-64.
4. Suetta C, Frandsen U, Mackey AL, Jensen L, Hvid LG, Bayer ML, Petersson SJ, Schrøder H, Andersen JL, Aagaard P, Schjerling P, Kjaer M. Ageing is associated with diminished muscle re-growth and myogenic precursor cell expansion early after immobility-induced atrophy in human skeletal muscle. *The Journal of Physiology*. 2013;591(Pt 15):3789-804.
5. Suetta C, Hvid LG, Justesen L, Christensen U, Neergaard K, Simonsen L, Ortenblad N, Magnusson SP, Kjaer M, Aagaard P. Effects of aging on human skeletal muscle after immobilization and retraining. *Journal of Applied Physiology*. 2009;107(4):1172-80.
6. Jensen C, Aagaard P, Overgaard S. Recovery in mechanical muscle strength following resurfacing vs standard total hip arthroplasty - a randomised clinical trial. *Osteoarthritis and cartilage* 2011;19(9):1108-16.
7. Jorgensen JR, Bech-Pedersen DT, Zeeman P, Sorensen J, Andersen LL, Schonberger M. Effect of intensive outpatient physical training on gait performance and cardiovascular health in people with hemiparesis after stroke. *Physical Therapy*. 2010;90(4):527-37.
8. Andersen LL, Zeeman P, Jorgensen JR, Bech-Pedersen DT, Sorensen J, Kjaer M, Andersen JL. Effects of intensive physical rehabilitation on neuromuscular adaptations in adults with poststroke hemiparesis. *Journal of strength and conditioning research* 2011; 25(10): 2808-17.
9. Liu-Ambrose TY, Khan KM, Eng JJ, Heinonen A, McKay HA. Both resistance and agility training increase cortical bone density in 75- to 85-year-old women with low bone mass: a 6-month randomized controlled trial. *Journal of clinical densitometry : the official journal of the International Society for Clinical Densitometry*. 2004;7(4):390-8.
10. Strasser B, Leeb G, Strehlow C, Schobersberger W, Haber P, Cauza E. The effects of strength and endurance training in patients with rheumatoid arthritis. *Clinical Rheumatology*. 2011;30(5):623-32.
11. Lemmey AB, Williams SL, Marcora SM, Jones J, Maddison PJ. Are the benefits of a high-intensity progressive resistance training program sustained in rheumatoid arthritis patients? A 3-year followup study. *Arthritis Care & Research*. 2012;64(1):71-5.
12. Ortega F, Toral J, Cejudo P, Villagomez R, Sanchez H, Castillo J, et al. Comparison of effects of strength and endurance training in patients with chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine*. 2002;166(5):669-74.
13. Kongsgaard M, Backer V, Jorgensen K, Kjaer M, Beyer N. Heavy resistance training increases muscle size, strength and physical function in elderly male COPD-patients--a pilot study. *Respiratory Medicine*. 2004;98(10):1000-7.

14. Aalund PK, Larsen K, Hansen TB, Bandholm T. Normalized knee-extension strength or leg-press power after fast-track total knee arthroplasty: which measure is most closely associated with performance-based and self-reported function? *Archives of Physical Medicine and Rehabilitation*. 2013;94(2):384-90.
15. Andersen LL, Kjaer M, Sogaard K, Hansen L, Kryger AI, Sjogaard G. Effect of two contrasting types of physical exercise on chronic neck muscle pain. *Arthritis and Rheumatism*. 2008;59(1):84-91.
16. Sjogaard G, Zebis MK, Kiilerich K, Saltin B, Pilegaard H. Exercise training and work task induced metabolic and stress-related mRNA and protein responses in myalgic muscles. *BioMed Research International*. 2013;2013:984523.
17. Langberg H, Ellingsgaard H, Madsen T, Jansson J, Magnusson SP, Aagaard P, Kjaer M. Eccentric rehabilitation exercise increases peritendinous type I collagen synthesis in humans with Achilles tendinosis. *Scandinavian Journal of Medicine & Science in Sports*. 2007;17(1):61-6.
18. Kongsgaard M, Kovanen V, Aagaard P, Doessing S, Hansen P, Laursen AH, Kjaer M, Magnusson SP. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scandinavian Journal of Medicine & Science in Sports*. 2009;19(6):790-802.
19. Vanhees L, Rauch B, Piepoli M, van Buuren F, Takken T, Borjesson M, et al. Importance of characteristics and modalities of physical activity and exercise in the management of cardiovascular health in individuals with cardiovascular disease (Part III). *European Journal of Preventive Cardiology*. 2012;19(6):1333-56.
20. Beyer N, Simonsen L, Bulow J, Lorenzen T, Jensen DV, Larsen L, Rasmussen U, Rennie M, Kjaer M. Old women with a recent fall history show improved muscle strength and function sustained for six months after finishing training. *Aging Clinical and Experimental Research*. 2007;19(4):300-9.
21. Nagarajan K, Bennett A, Agostini P, Naidu B. Is preoperative physiotherapy/pulmonary rehabilitation beneficial in lung resection patients? *Interactive Cardiovascular and Thoracic Surgery*. 2011;13(3):300-2.
22. Handoll HH, Sherrington C, Mak JC. Interventions for improving mobility after hip fracture surgery in adults. *The Cochrane Database of Systematic Reviews*. 2011(3):CD001704.
23. Kronborg L, Bandholm T, Palm H, Kehlet H, Kristensen MT. Feasibility of progressive strength training implemented in the acute ward after hip fracture surgery. *PloS one*. 2014;9(4):e93332.
24. Alkjaer T, Wieland MR, Andersen MS, Simonsen EB, Rasmussen J. Computational modeling of a forward lunge: towards a better understanding of the function of the cruciate ligaments. *Journal of Anatomy*. 2012;221(6):590-7.
25. Aagaard P, Simonsen EB, Andersen JL, Magnusson P, Dyhre-Poulsen P. Increased rate of force development and neural drive of human skeletal muscle following resistance training. *Journal of Applied Physiology*. 2002;93(4):1318-26.
26. Bojsen-Moller J, Schwartz S, Kalliokoski KK, Finni T, Magnusson SP. Intermuscular force transmission between human plantarflexor muscles in vivo. *Journal of Applied Physiology*. 2010;109(6):1608-18.
27. Magnusson SP, Hansen P, Aagaard P, Brond J, Dyhre-Poulsen P, Bojsen-Moller J, Kjaer M. Differential strain patterns of the human gastrocnemius aponeurosis and free tendon, in vivo. *Acta Physiologica Scandinavica*. 2003;177(2):185-95.
28. Cronin NJ, Lichtwark G. The use of ultrasound to study muscle-tendon function in human posture and locomotion. *Gait & Posture*. 2013;37(3):305-12.
29. Bojsen-Moller J, Losnegard T, Kemppainen J, Viljanen T, Kalliokoski KK, Hallen J. Muscle use during double poling evaluated by positron emission tomography. *Journal of Applied Physiology*. 2010;109(6):1895-903.
30. Masood T, Bojsen-Moller J, Kalliokoski KK, Kirjavainen A, Aarimaa V, Peter Magnusson S, et al. Differential contributions of ankle plantarflexors during submaximal isometric muscle action: a PET and EMG study. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology*. 2014;24(3):367-74.
31. Bojsen-Moller J, Kalliokoski K, Seppanen M, Kjaer M, Magnusson SP. Low-intensity tensile loading increases intratendinous glucose uptake in the Achilles tendon. *Journal of Applied Physiology*. 2006;101(1):196-201.

32. Enocson AG, Berg HE, Vargas R, Jenner G, Tesch PA. Signal intensity of MR-images of thigh muscles following acute open- and closed chain kinetic knee extensor exercise - index of muscle use. *European Journal of Applied Physiology*. 2005;94(4):357-63.
33. Finni T, Hodgson JA, Lai AM, Edgerton VR, Sinha S. Mapping of movement in the isometrically contracting human soleus muscle reveals details of its structural and functional complexity. *Journal of Applied Physiology*. 2003;95(5):2128-33.
34. Fleckenstein JL, Canby RC, Parkey RW, Peshock RM. Acute effects of exercise on MR imaging of skeletal muscle in normal volunteers. *American journal of Roentgenology*. 1988;151(2):231-7.
35. Andersen LL, Magnusson SP, Nielsen M, Haleem J, Poulsen K, Aagaard P. Neuromuscular activation in conventional therapeutic exercises and heavy resistance exercises: implications for rehabilitation. *Physical therapy*. 2006;86(5):683-97.
36. Wilk KE, Escamilla RF, Fleisig GS, Barrentine SW, Andrews JR, Boyd ML. A comparison of tibiofemoral joint forces and electromyographic activity during open and closed kinetic chain exercises. *The American Journal of Sports Medicine*. 1996;24(4):518-27.
37. Aagaard P, Simonsen EB, Andersen JL, Magnusson SP, Bojsen-Moller F, Dyhre-Poulsen P. Antagonist muscle coactivation during isokinetic knee extension. *Scandinavian Journal of Medicine & Science in Sports*. 2000;10(2):58-67.
38. Zebis MK, Andersen LL, Bencke J, Kjaer M, Aagaard P. Identification of athletes at future risk of anterior cruciate ligament ruptures by neuromuscular screening. *The American Journal of Sports Medicine*. 2009;37(10):1967-73.
39. Escamilla RF, Fleisig GS, Zheng N, Lander JE, Barrentine SW, Andrews JR, et al. Effects of technique variations on knee biomechanics during the squat and leg press. *Medicine and Science in Sports and Exercise*. 2001;33(9):1552-66.
40. Jakobsen MD, Sundstrup E, Andersen CH, Aagaard P, Andersen LL. Muscle activity during leg strengthening exercise using free weights and elastic resistance: effects of ballistic vs controlled contractions. *Human movement science*. 2013;32(1):65-78.
41. Heijne A, Fleming BC, Renstrom PA, Peura GD, Beynon BD, Werner S. Strain on the anterior cruciate ligament during closed kinetic chain exercises. *Medicine and Science in Sports and Exercise*. 2004;36(6):935-41.
42. Suetta C, Aagaard P, Magnusson SP, Andersen LL, Sipila S, Rosted A, Jakobsen AK, Duus B, Kjaer M. Muscle size, neuromuscular activation, and rapid force characteristics in elderly men and women: effects of unilateral long-term disuse due to hip-osteoarthritis. *Journal of Applied Physiology*. 2007;102(3):942-8.
43. Maffiuletti NA, Minetto MA, Farina D, Bottinelli R. Electrical stimulation for neuromuscular testing and training: state-of-the art and unresolved issues. *European Journal of Applied Physiology*. 2011;111(10):2391-7.
44. Kim KM, Croy T, Hertel J, Saliba S. Effects of neuromuscular electrical stimulation after anterior cruciate ligament reconstruction on quadriceps strength, function, and patient-oriented outcomes: a systematic review. *The Journal of Orthopaedic and Sports Physical Therapy*. 2010;40(7):383-91.
45. Melo MD, Pompeo KD, Brodt GA, Baroni BM, da Silva Junior DP, Vaz MA. Effects of neuromuscular electrical stimulation and low-level laser therapy on the muscle architecture and functional capacity in elderly patients with knee osteoarthritis: a randomized controlled trial. *Clinical Rehabilitation*. In press 2014
46. Elboim-Gabyzon M, Rozen N, Laufer Y. Does neuromuscular electrical stimulation enhance the effectiveness of an exercise programme in subjects with knee osteoarthritis? A randomized controlled trial. *Clinical Rehabilitation*. 2013;27(3):246-57.
47. Giggins O, Fullen B, Coughlan G. Neuromuscular electrical stimulation in the treatment of knee osteoarthritis: a systematic review and meta-analysis. *Clinical Rehabilitation*. 2012;26(10):867-81.
48. Maddocks M, Gao W, Higginson IJ, Wilcock A. Neuromuscular electrical stimulation for muscle weakness in adults with advanced disease. *The Cochrane database of systematic reviews*. 2013;1:CD009419.
49. Williams N, Flynn M. A review of the efficacy of neuromuscular electrical stimulation in critically ill patients. *Physiotherapy Theory and Practice*. 2014;30(1):6-11.

50. Feil S, Newell J, Minogue C, Paessler HH. The effectiveness of supplementing a standard rehabilitation program with superimposed neuromuscular electrical stimulation after anterior cruciate ligament reconstruction: a prospective, randomized, single-blind study. *The American Journal of Sports Medicine*. 2011;39(6):1238-47.
51. Maffiuletti NA, Vivodtzev I, Minetto MA, Place N. A new paradigm of neuromuscular electrical stimulation for the quadriceps femoris muscle. *European Journal of Applied Physiology*. 2014;114(6):1197-205.
52. Wellauer V, Morf C, Minetto MA, Place N, Maffiuletti NA. Assessment of quadriceps muscle inactivation with a new electrical stimulation paradigm. *Muscle & Nerve*. 2014.
53. Manini TM, Clark BC. Blood flow restricted exercise and skeletal muscle health. *Exercise and Sport Sciences Reviews*. 2009;37(2):78-85.
54. Nielsen JL, Aagaard P, Bech RD, Nygaard T, Hvid LG, Wernbom M, Suetta C, Frandsen U. Proliferation of myogenic stem cells in human skeletal muscle in response to low-load resistance training with blood flow restriction. *The Journal of Physiology*. 2012;590(Pt 17):4351-61.
55. Abe T, Kearns CF, Sato Y. Muscle size and strength are increased following walk training with restricted venous blood flow from the leg muscle, Kaatsu-walk training. *Journal of Applied Physiology*. 2006;100(5):1460-6.
56. Takarada Y, Takazawa H, Sato Y, Takebayashi S, Tanaka Y, Ishii N. Effects of resistance exercise combined with moderate vascular occlusion on muscular function in humans. *Journal of Applied Physiology*. 2000;88(6):2097-106.
57. Loenneke JP, Abe T, Wilson JM, Thiebaud RS, Fahs CA, Rossow LM, et al. Blood flow restriction: an evidence based progressive model (Review). *Acta Physiologica Hungarica*. 2012;99(3):235-50.
58. Fry CS, Glynn EL, Drummond MJ, Timmerman KL, Fujita S, Abe T, et al. Blood flow restriction exercise stimulates mTORC1 signaling and muscle protein synthesis in older men. *Journal of Applied Physiology*. 2010;108(5):1199-209.
59. Mikkelsen LR, Mechlenburg I, Søballe K, Jørgensen LB, Mikkelsen S, Bandholm T, Petersen AK. Effect of early supervised progressive resistance training compared to unsupervised home-based exercise after fast-track total hip replacement applied to patients with preoperative functional limitations. A single-blinded randomised controlled trial. *Osteoarthritis and Cartilage*, in press 2014
60. Zebis MK, Skotte J, Andersen CH, Mortensen P, Petersen MH, Viskær TC, Jensen TL, Bencke J, Andersen LL. Kettlebell swing targets semitendinosus and supine leg curl targets biceps femoris: an EMG study with rehabilitation implications. *British Journal of Sports Medicine*. 2013;47:1192-1198.