

Referencer til artikel i Dansk Sportsmedicin nr. 4, 2014:

## Styrketræning ved kronisk seneoverbelastning

Af Christian Couppé<sup>1,2</sup> og Jens Olesen<sup>1</sup>

<sup>1</sup>Institut for Idrætsmedicin, København og <sup>2</sup>Forskningsenheden for Muskuloskeletal Rehabilitering, Fysio- og ergoterapiafdelingen, Bispebjerg og Frederiksberg Hospitaler

- ALFREDSON, H. 2005. The chronic painful Achilles and patellar tendon: research on basic biology and treatment. *Scand J Med Sci Sports*, 15, 252-259.
- ANDERSEN, L. L., MAGNUSSON, S. P., NIELSEN, M., HALEEM, J., POULSEN, K. & AAGAARD, P. 2006. Neuromuscular activation in conventional therapeutic exercises and heavy resistance exercises: implications for rehabilitation. *Phys Ther*, 86, 683-697.
- ARNOCZKY, S. P., LAVAGNINO, M. & EGERBACHER, M. 2007. The mechanobiological aetiopathogenesis of tendinopathy: is it the over-stimulation or the under-stimulation of tendon cells? *Int.J.Exp.Pathol.*, 88, 217-226.
- COUPPE, C., KONGSGAARD, M., AAGAARD, P., HANSEN, P., BOJSEN-MOLLER, J., KJAER, M. & MAGNUSSON, S. P. 2008. Habitual loading results in tendon hypertrophy and increased stiffness of the human patellar tendon. *J Appl Physiol*, 105, 805-10.
- COUPPE, C., KONGSGAARD, M., AAGAARD, P., VINTHER, A., BOESEN, M., KJAER, M. & MAGNUSSON, S. P. 2013. Differences in tendon properties in elite badminton players with or without patellar tendinopathy. *Scand J Med Sci Sports*, 23, e89-95.
- FARUP, J., RAHBEK, S. K., VENDELBO, M. H., MATZON, A., HINDHEDE, J., BEJDER, A., RINGGARD, S. & VISSING, K. 2013. Whey protein hydrolysate augments tendon and muscle hypertrophy independent of resistance exercise contraction mode. *Scand J Med Sci Sports*.
- FROHM, A., HALVORSEN, K. & THORSTENSSON, A. 2007a. Patellar tendon load in different types of eccentric squats. *Clin.Biomech.(Bristol, Avon.)*, 22, 704-711.
- FROHM, A., SAARTOK, T., HALVORSEN, K. & RENSTROM, P. 2007b. Eccentric treatment for patellar tendinopathy: a prospective randomised short-term pilot study of two rehabilitation protocols. *Br.J Sports Med.*, 41, e7.
- JONSSON, P. & ALFREDSON, H. 2005. Superior results with eccentric compared to concentric quadriceps training in patients with jumper's knee: a prospective randomised study. *Br J Sports Med*, 39, 847-50.
- KHAN, K. M., COOK, J. L., BONAR, F., HARCOURT, P. & ASTROM, M. 1999. Histopathology of common tendinopathies. Update and implications for clinical management. *Sports Med*, 27, 393-408.
- KJAER, M. & HEINEMEIER, K. M. 2014. 4. Eccentric exercise: Acute and chronic effects on healthy and diseased tendons. *J Appl Physiol (1985)*.
- KONGSGAARD, M., KOVANEN, V., AAGAARD, P., DOESSING, S., HANSEN, P., LAURSEN, A. H., KALDAU, N. C., KJAER, M. & MAGNUSSON, S. P. 2009. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scand.J.Med.Sci.Sports*, 19, 790-802.
- KONGSGAARD, M., QVORTRUP, K., LARSEN, J., AAGAARD, P., DOESSING, S., HANSEN, P., KJAER, M. & MAGNUSSON, S. P. 2010. Fibril morphology and tendon mechanical properties in

- patellar tendinopathy: effects of heavy slow resistance training. *Am.J.Sports Med.*, 38, 749-756.
- KONGSGAARD, M., REITELSEDER, S., PEDERSEN, T. G., HOLM, L., AAGAARD, P., KJAER, M. & MAGNUSSON, S. P. 2007. Region specific patellar tendon hypertrophy in humans following resistance training. *Acta Physiol (Oxf)*, 191, 111-121.
- KONGSGAARD, M., AAGAARD, P., ROIKJAER, S., OLSEN, D., JENSEN, M., LANGBERG, H. & MAGNUSSON, S. P. 2006. Decline eccentric squats increases patellar tendon loading compared to standard eccentric squats. *Clin.Biomech.(Bristol, Avon.)*, 21, 748-754.
- MAEDA, E., SHELTON, J. C., BADER, D. L. & LEE, D. A. 2007. Time dependence of cyclic tensile strain on collagen production in tendon fascicles. *Biochem.Biophys.Res.Commun.*, 362, 399-404.
- MAFFULLI, N., TESTA, V., CAPASSO, G., EWEN, S. W., SULLO, A., BENAZZO, F. & KING, J. B. 2004. Similar histopathological picture in males with Achilles and patellar tendinopathy. *Med.Sci.Sports Exerc.*, 36, 1470-1475.
- MAGNUSSON, S. P., LANGBERG, H. & KJAER, M. 2010. The pathogenesis of tendinopathy: balancing the response to loading. *Nat.Rev.Rheumatol.*, 6, 262-268.
- MALLIARAS, P., BARTON, C. J., REEVES, N. D. & LANGBERG, H. 2013. Achilles and patellar tendinopathy loading programmes : a systematic review comparing clinical outcomes and identifying potential mechanisms for effectiveness. *Sports Med*, 43, 267-86.
- RATHLEFF, M. S., MOLGAARD, C. M., FREDBERG, U., KAALUND, S., ANDERSEN, K. B., JENSEN, T. T., AASKOV, S. & OLESEN, J. L. 2014. High-load strength training improves outcome in patients with plantar fasciitis: A randomized controlled trial with 12-month follow-up. *Scand J Med Sci Sports*.
- SEYNNES, O. R., ERSKINE, R. M., MAGANARIS, C. N., LONGO, S., SIMONEAU, E. M., GROSSET, J. F. & NARICI, M. V. 2009. Training-induced changes in structural and mechanical properties of the patellar tendon are related to muscle hypertrophy but not to strength gains. *J Appl Physiol*, 107, 523-30.
- SILBERNAGEL, K. G., THOMEE, R., ERIKSSON, B. I. & KARLSSON, J. 2007. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled study. *Am J Sports Med*, 35, 897-906.
- VISNES, H., HOKSRUD, A., COOK, J. & BAHR, R. 2005. No effect of eccentric training on jumper's knee in volleyball players during the competitive season: a randomized clinical trial. *Clin.J Sport Med*, 15, 227-234.
- WASIELEWSKI, N. J. & KOTSKO, K. M. 2007. Does eccentric exercise reduce pain and improve strength in physically active adults with symptomatic lower extremity tendinosis? A systematic review. *J Athl Train*, 42, 409-21.
- WOODLEY, B. L., NEWSHAM-WEST, R. J. & BAXTER, G. D. 2007. Chronic tendinopathy: effectiveness of eccentric exercise. *Br.J Sports Med.*, 41, 188-198.
- YOUNG, M. A., COOK, J. L., PURDAM, C. R., KISS, Z. S. & ALFREDSON, H. 2005. Eccentric decline squat protocol offers superior results at 12 months compared with traditional eccentric protocol for patellar tendinopathy in volleyball players. *Br J Sports Med*, 39, 102-5.