

**References** for the article in Dansk Sportsmedicin nr. 1, 2015:

## “Exercise as treatment for the middle-aged presenting with a meniscal tear”

*Ewa M Roos<sup>1</sup> and May Arna Risberg<sup>2</sup>*

<sup>1</sup>*Research Unit for Musculoskeletal Function and Physiotherapy, Department of Sports and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark*

<sup>2</sup>*Norwegian research center for Active Rehabilitation, Department of Orthopedic Surgery, Oslo University Hospital, Oslo, Norway*

1. Englund M, Roemer FW, Hayashi D, Crema MD, Guermazi A. Meniscus pathology, osteoarthritis and the treatment controversy. *Nat Rev Rheumatol.* 2012;8(7):412-9.
2. Thorlund JB, Hare KB, Lohmander LS. Large increase in arthroscopic meniscus surgery in the middle-aged and older population in Denmark from 2000 to 2011. *Acta Orthop.* 2014;85(3):287-92.
3. Oiestad BE, Engebretsen L, Storheim K, Risberg MA. Knee osteoarthritis after anterior cruciate ligament injury: a systematic review. *Am J Sports Med.* 2009;37(7):1434-43.
4. Oiestad BE, Holm I, Gunderson R, Myklebust G, Risberg MA. Quadriceps muscle weakness after anterior cruciate ligament reconstruction: a risk factor for knee osteoarthritis? *Arthritis Care Res (Hoboken).* 2010;62(12):1706-14.
5. Felson DT, Lawrence RC, Dieppe PA, Hirsch R, Helmick CG, Jordan JM, et al. Osteoarthritis: new insights. Part 1: the disease and its risk factors. *Ann Intern Med.* 2000;133(8):635-46.
6. Seon JK, Song EK, Park SJ. Osteoarthritis after anterior cruciate ligament reconstruction using a patellar tendon autograft. *Int Orthop.* 2006;30(2):94-8.
7. Stensrud S, Risberg MA, Roos EM. Knee function and knee muscle strength in middle-aged patients with degenerative meniscal tears eligible for arthroscopic partial meniscectomy. *British journal of sports medicine.* 2012.
8. Peat G, Croft P, Hay E. Clinical assessment of the osteoarthritis patient. *Best Pract Res Clin Rheumatol.* 2001;15(4):527-44.
9. Kittelson AJ, Thomas AC, Kluger BM, Stevens-Lapsley JE. Corticospinal and intracortical excitability of the quadriceps in patients with knee osteoarthritis. *Exp Brain Res.* 2014;232(12):3991-9.
10. McAlindon TE, Cooper C, Kirwan JR, Dieppe PA. Determinants of disability in osteoarthritis of the knee. *Ann Rheum Dis.* 1993;52(4):258-62.
11. Liikavainio T, Isolehto J, Helminen HJ, Perttunen J, Lepola V, Kiviranta I, et al. Loading and gait symmetry during level and stair walking in asymptomatic subjects with knee osteoarthritis: importance of quadriceps femoris in reducing impact force during heel strike? *Knee.* 2007;14(3):231-8.
12. Øiestad BE, Juhl CB, Eitzen I, Thorlund JB. Knee extensor muscle weakness is a risk factor for development of knee osteoarthritis. A systematic review and meta-analysis. *Osteoarthritis and Cartilage.* 2014;In revision.

13. Mikesky AE, Meyer A, Thompson KL. Relationship between quadriceps strength and rate of loading during gait in women. *J Orthop Res.* 2000;18(2):171-5.
14. Bennell KL, Hunt MA, Wrigley TV, Hunter DJ, McManus FJ, Hodges PW, et al. Hip strengthening reduces symptoms but not knee load in people with medial knee osteoarthritis and varus malalignment: a randomised controlled trial. *Osteoarthritis Cartilage.* 2010;18(5):621-8.
15. Ageberg E, Roos EM. Neuromuscular exercise as treatment of degenerative knee disease. *Exercise and sport sciences reviews.* 2014.
16. Bennell KL, Wrigley TV, Hunt MA, Lim BW, Hinman RS. Update on the role of muscle in the genesis and management of knee osteoarthritis. *Rheum Dis Clin North Am.* 2013;39(1):145-76.
17. Knæartrose – nationale kliniske retningslinjer og faglige visitationsretningslinjer. København: Sundhedsstyrelsen, 2012.
18. McAlindon TE, Bannuru RR, Sullivan MC, Arden NK, Berenbaum F, Bierma-Zeinstra SM, et al. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis and cartilage / OARS, Osteoarthritis Research Society.* 2014;22(3):363-88.
19. Messier SP, Gutekunst DJ, Davis C, DeVita P. Weight loss reduces knee-joint loads in overweight and obese older adults with knee osteoarthritis. *Arthritis Rheum.* 2005;52(7):2026-32.
20. Wang SY, Olson-Kellogg B, Shamliyan TA, Choi JY, Ramakrishnan R, Kane RL. Physical therapy interventions for knee pain secondary to osteoarthritis: a systematic review. *Ann Intern Med.* 2012;157(9):632-44.
21. Juhl C, Christensen R, Roos EM, Zhang W, Lund H. Impact of exercise type and dose on pain and disability in knee osteoarthritis: a systematic review and meta-regression analysis of randomized controlled trials. *Arthritis Rheumatol.* 2014;66(3):622-36.
22. Stensrud S, Roos EM, Risberg MA. A 12-Week Exercise Therapy Program in Middle-Aged Patients With Degenerative Meniscus Tears: A Case Series With 1-Year Follow-up. *The Journal of orthopaedic and sports physical therapy.* 2012;42(11):919-31.
23. Ageberg E, Link A, Roos EM. Feasibility of neuromuscular training in patients with severe hip or knee OA: the individualized goal-based NEMEX-TJR training program. *BMC musculoskeletal disorders.* 2010;11:126.
24. Cotofana S, Ring-Dimitriou S, Hudelmaier M, Himmer M, Wirth W, Sanger AM, et al. Effects of exercise intervention on knee morphology in middle-aged women: a longitudinal analysis using magnetic resonance imaging. *Cells, tissues, organs.* 2010;192(1):64-72.
25. Hovis KK, Stehling C, Souza RB, Haughom BD, Baum T, Nevitt M, et al. Physical activity is associated with magnetic resonance imaging-based knee cartilage T2 measurements in asymptomatic subjects with and those without osteoarthritis risk factors. *Arthritis Rheum.* 2011;63(8):2248-56.
26. Van Ginckel A, Witvrouw E. Acute cartilage loading responses after an in vivo squatting exercise in people with doubtful to mild knee osteoarthritis: a case-control study. *Phys Ther.* 2013;93(8):1049-60.
27. Steib S, Schoene D, Pfeifer K. Dose-response relationship of resistance training in older adults: a meta-analysis. *Med Sci Sports Exerc.* 2010;42(5):902-14.
28. Khan M, Evaniew N, Bedi A, Ayeni OR, Bhandari M. Arthroscopic surgery for degenerative tears of the meniscus: a systematic review and meta-analysis. *CMAJ.* 2014;186(14):1057-64.

29. Osteras H, Osteras B, Torstensen TA. Medical exercise therapy, and not arthroscopic surgery, resulted in decreased depression and anxiety in patients with degenerative meniscus injury. *Journal of bodywork and movement therapies*. 2012;16(4):456-63.
30. Stensrud S, Risberg MA, Roos EM. Effect of Exercise Therapy Compared with Arthroscopic Surgery on Knee Muscle Strength and Functional Performance in Middle-Aged Patients with Degenerative Meniscus Tears: A 3-Mo Follow-up of a Randomized Controlled Trial. *Am J Phys Med Rehabil*. 2014.
31. Bedi A, Kelly NH, Baad M, Fox AJ, Brophy RH, Warren RF, et al. Dynamic contact mechanics of the medial meniscus as a function of radial tear, repair, and partial meniscectomy. *J Bone Joint Surg Am*. 2010;92(6):1398-408.
32. Lohmander LS, Englund PM, Dahl LL, Roos EM. The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *Am J Sports Med*. 2007;35(10):1756-69.
33. Roos EM, Dahlberg L. Positive effects of moderate exercise on knee cartilage glycosaminoglycan content. A four-month randomized controlled trial in patients at risk of osteoarthritis. *Arthritis Rheum*. 2005;52:3507-14.
34. Owman H, Ericsson YB, Englund M, Tiderius CJ, Tjornstrand J, Roos EM, et al. Association between delayed gadolinium-enhanced MRI of cartilage (dGEMRIC) and joint space narrowing and osteophytes: a cohort study in patients with partial meniscectomy with 11 years of follow-up. *Osteoarthritis Cartilage*. 2014;22(10):1537-41.