

Referencer til artikel i Dansk Sportsmedicin nr. 1, 2014:

High Volume Injection (HVI) Behandling af kroniske seneproblemer

Af Morten Boesen, speciallæge i ortopædkirurgi, Teres Parken Privathospital; Henning Langberg, professor; Anders Boesen, læge, PhD-stud, Idrætsmedicinsk Institut, Bispebjerg Hospital.

- 1) Longo, U. G., Rittweger, J., Garau, G., Radonic, B., Gutwasser, C., Gilliver, S. F., et al. (2009). No influence of age, gender, weight, height, and impact profile in achilles tendinopathy in masters track and field athletes. *The American Journal of Sports Medicine*, 37(7), 1400e1405.
- 2) Alfredson H, Lorentzon R. Chronic Achilles tendinosis: recommendations for treatment and prevention. *Sports Med* 2000. 29:135–146.
- 3) Schepsis A A, Jones H, Haas A L. Achilles tendon disorders in athletes. *Am J Sports Med* 2002. 30:287–305.
- 4) Kujala UM, Sarna S, Kaprio J. Cumulative incidence of Achilles tendon rupture and tendinopathy in male former elite athletes. *Clin J Sport Med* 2005;15(3):133 – 135.
- 5) Alfredson H. Chronic midportion Achilles tendinopathy: an update on research and treatment. *Clin Sports Med* 2003;22: 727–41
- 6) Astrom M. Partial rupture in chronic Achilles tendinopathy. A retrospective analysis of 342 cases. *Acta Orthop Scand* 1998; 69:404–7
- 7) Luscombe KL, Sharma P, Maffulli N. Achilles tendinopathy. *Trauma* 2003;5:215–25
- 8) Rolf C, Movin T. Etiology, histopathology, and outcome of surgery in achillodynia. *Foot Ankle Int* 1997;18:565–9
- 9) Cook J L, Khan K M, Purdam C. Achilles tendinopathy. *Man Ther* 2002. 7:121–130.
- 10) de Jonge S, et al. Incidence of midportion Achilles tendinopathy in the general population. *Br J Sports Med* 2011; 45: 1026–8. [
- 11) Järvinen TAH, et al. Achilles tendon disorders: etiology and epidemiology. *Foot Ankle Clin N Am* 2005; 10: 255–66.
- 12) Carcia CR, et al. Achilles pain, stiffness, and muscle power deficits: Achilles teninitis. *J Orthop Sport Phys Ther* 2010; 40: A1–26.
- 13) Holmes GB, Lin J. Etiologic factors associated with symptomatic Achilles tendinopathy. *Foot Ankle Int* 2006; 27: 952–9.
- 14) Klemp P, et al. Musculoskeletal manifestations in hyperlipidaemia: a controlled study. *Ann Rheum Dis* 1993; 52: 44–8.
- 15) Doyle HE. Tendinopathy resulting from the use of fluoroquinolones: managing risks. *JAAPA* 2010; 23: 18–21, 54.
- 16) Mahieu NN, Witvrouw E, Stevens V, Van TD, Roget P. Intrinsic risk factors for the development of Achilles tendon overuse injury: A prospective study. *Am J Sports Med* 2006;34(2):226 – 235.
- 17) Kvist M. Achilles tendon injuries in athletes. *Ann Chir Gynaecol* 1991;80(2):188 – 201

- 18) Ames, P. R., Longo, U. G., Denaro, V., & Maffulli, N. (2008). Achilles tendon problems: not just an orthopaedic issue. *Disability and Rehabilitation*, 30(20e22), 1646e1650.
- 19) Longo, U. G., Oliva, F., Denaro, V., & Maffulli, N. (2008). Oxygen species and overuse tendinopathy in athletes. *Disability and Rehabilitation*, 30, 20e22, 1563e1571.
- 20) Battery L, Maffulli N. Inflammation in overuse tendon injuries. *Sports Med Arthrosc Rev* 2011; 19: 213–7.
- 21) Maffulli N, *et al.* Current treatment options for tendinopathy. *Expert Opin Pharmacother* 2010; 11: 2177–86.
- 22) Knobloch K, Kraemer R, Lichtenberg A, Jagodzinski M, Gossling T, Richter M, *et al.* Achilles tendon and paratendon microcirculation in midportion and insertional tendinopathy in athletes. *Am J Sports Med* 2006;34(1):92 – 97.
- 23) Alfredson H, Ohberg L, Forsgren S. Is vasculo-neural ingrowth the cause of pain in chronic Achilles tendinosis? An investigation using ultrasonography and colour Doppler, immunohistochemistry, and diagnostic injections. *Knee Surg Sports Traumatol Arthrosc* 2003;11(5):334 – 338.
- 24) Kristoffersen M, Ohberg L, Johnston C, Alfredson H. Neovascularisation in chronic tendon injuries detected with colour Doppler ultrasound in horse and man: implications for research and treatment. *Knee Surg Sports Traumatol Arthrosc* 2005;13(6):505 – 508
- 25) Ohberg L, Lorentzon R, Alfredson H. Neovascularisation in Achilles tendons with painful tendinosis but not in normal tendons: An ultrasonographic investigation. *Knee Surg Sports Traumatol Arthrosc* 2001;9(4):233 – 238
- 26) Longo, U. G., Ronga, M., & Maffulli, N. (2009). Achilles tendinopathy. *Sports Medicine and Arthroscopy Review*, 17(2), 112e126.
- 27) Ohberg L, Lorentzon R, Alfredson H. Neovascularisation in Achilles tendons with painful tendinosis but not in normal tendons: an ultrasonographic investigation. *Knee Surg Sports Traumatol Arthrosc* 2001;9:233–8.
- 28) Aström M, Rausing A. Chronic Achilles tendinopathy. A survey of surgical and histopathologic findings. *Clin Orthop Relat Res* 1995;316:151–64.
- 29) Jozsa L, Kannus P. Human tendons: anatomy, physiology, path- ology. Champaign, IL: Human Kinetics; 1997. p. 1–574.
- 30) Scott A, *et al.* Conservative treatment of chronic Achilles tendinopathy. *CMAJ* 2011; 183: 1159–65.
- 31) Magnussen RA, *et al.* Nonoperative treatment of midportion Achilles tendinopathy: a systematic review. *Clin J Sport Med* 2009; 19: 54–64.
- 32) Silbernagel KG, *et al.* Eccentric overload training for patients with chronic Achilles tendon pain—a randomised controlled study with reliability testing of the evaluation methods. *Scand J Med Sci Sports* 2001; 11: 197–206
- 33) ArchambaultJM,WileyJP,BrayRC,VerhoefM,WisemanDA, Elliott PD. Can sonography predict the outcome in patients with achillodynia? *J Clin Ultrasound* 1998;26(7):335–9.

- 34) Hartgerink P, Fessell DP, Jacobson JA, van Holsbeeck MT. Full- versus partial-thickness Achilles, tendon tears: sonographic accuracy and characterization in 26 cases with surgical correlation. *Radiology* 2001;220:406–12.
- 35) Öhberg L, Lorentzon R, Alfredson H. Neovascularisation in Achilles tendons with painful tendinosis but not in normal tendons: an ultrasonographic investigation. *Knee Surg Sports Traumatol Arthrosc* 2001;9:233–8.*
- 36) Scott A, *et al.* Conservative treatment of chronic Achilles tendinopathy. *CMAJ* 2011; 183: 1159–65.
- 37) Alfredson H, Lorentzon R. Chronic Achilles tendinosis: recommendations for treatment and prevention. *Sports Med* 2000; 29:135–46.
- 38) Starish WD, Rubinovich RM, Curwin S. Eccentric exercise in chronic tendinitis. *Clin Orthop Rehab Res* 1986;208:65–8.
- 39) Rees JD, Wolman RC, Wilson A. Eccentric exercises; why do they work, what are the problems and how can we improve them? *Br J Sports Med* 2009;43:242–6.
- 40) Mayer F, *et al.* Effects of short-term treatment strategies over 4 weeks in Achilles tendinopathy. *Br J Sports Med* 2007; 41: e6.
- 41) Rompe JD, *et al.* Eccentric loading, shock-wave treatment, or a wait-and-see policy for tendinopathy of the main body of tendo Achillis: a randomized controlled trial. *Am J Sports Med* 2007; 35: 374–83.
- 42) Mafi N, *et al.* Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 2001; 9: 42–7.
- 43) Niesen-Vertommen SL, *et al.* The effect of eccentric versus concentric exercise in the management of Achilles tendonitis. *Clin J Sport Med* 1992; 2: 109–13.
- 44) Roos EM, *et al.* Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy—a randomized trial with 1-year follow-up. *Scand J Med Sci Sports* 2004; 14: 286–95.
- 45) Kingma JJ, de Knikker R, Wittink HM, Takken T. Eccentric overload training in patients with chronic Achilles tendinopathy: a systematic review. *Br J Sports Med* 2007;41. e3. Available from <http://www.bjsportmed.com/cgi/content/full/41/6/e3>.
- 46) Coombes BK, *et al.* Efficacy and safety of corticosteroid injection and other injections for management of tendinopathy: a systematic review of randomised controlled trials. *Lancet* 2010; 376: 1751–67.
- 47) Alfredson, H., & Ohberg, L. (2005). Sclerosing injections to areas of neo-vascularisation reduce pain in chronic Achilles tendinopathy: a double-blind randomised controlled trial.

- Knee Surgery, Sports Traumatology, Arthroscopy, 13(4), 338e344.
- 48) van Sterkenburg, M. N., de Jonge, M. C., Sierevelt, I. N., & van Dijk, C. N. (2010). Less promising results with sclerosing etoxysclerol injections for midportion Achilles tendinopathy: a retrospective study. *The American Journal of Sports Medicine*, [Epub ahead of print] Jul 2.
 - 49) National Institute for Health and Clinical Excellence, 2009. *Autologous blood injection for tendinopathy* [online]. Available: <http://www.nice.org.uk/nicemedia/pdf/IPG279Guidance.PDF>
 - 50) de Vos RJ, *et al.* Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. *JAMA* 2010; 303: 144–9.
 - 51) de Jonge S, *et al.* One-year follow-up of platelet-rich plasma treatment in chronic Achilles tendinopathy: a double-blind randomized placebo-controlled trial. *Am J Sports Med* 2011; 39: 1623–9.
 - 52) Ryan M, *et al.* Favorable outcomes after sonographically guided intratendinous injection of hyperosmolar dextrose for chronic insertional and midportion Achilles tendinosis. *AJR Am J Roentgenol* 2010; 194: 1047–53.
 - 53) Longo, U. G., Lamberti, A., Maffulli, N., & Denaro, V. (2010b). Tissue engineered biological augmentation for tendon healing: a systematic review. *British Medical Bulletin*, doi: 10.1093/bmb/ldq030 [pii] 10.1093/bmb/ldq030.
 - 54) Lippi, G., Longo, U. G., & Maffulli, N. (2010). Genetics and sports. *British Medical Bulletin*, 93(1), 27e47. <http://dx.doi.org/10.1093/Bmb/Ldp007>.
 - 55) Chan O, *et al.* High volume image guided injections in chronic Achilles tendinopathy. *Disabil Rehabil* 2008; 30: 1697–708.
 - 56) Humphrey, J., Chan, O., Crisp, T., Padhiar, N., Morrissey, D., Twycross-Lewis, R., et al. (2009). The short-term effects of high volume image guided injections in resistant non-insertional Achilles tendinopathy. *Journal of Sports Science and Medicine*,
 - 57) Resteghini, Peter; Yeoh, Justin. High-volume injection in the management of recalcitrant mid-body Achilles tendinopathy: a prospective case series assessing the influence of neovascularity and outcome. *International Musculoskeletal Medicine*, Volume 34, Number 3, September 2012 , pp. 92-100(9)
 - 58) Nicola Maffulli, Filippo Spiezia, Umile Giuseppe Longo, Vincenzo Denaro, Gayle D. Maffulli. High volume image guided injections for the management of chronic tendinopathy of the main body of the Achilles tendon. *Physical Therapy in Sport* xxx (2012) 1e5
 - 59) Hatim Abdulhusein, Dylan Morrissey, Otto Chan. High Volume Image Guided Injections with or without steroid for Achilles Tendinopathy: A Prospective Study, BSc 090124310, July 2013
 - 60) Johal, K. S.; Longman, C.; Malhi, A.; Madhav, R.; De Villiers, P. High volume injection for Achilles tendinopathy. *Muscles, Ligaments & Tendons Journal (MLTJ)*;2012 Supplement, p3
 - 61) Andren L, Lundberg B. Treatment of rigid shoulders by joint distension during arthrography. *Acta Orthop Scand* 1965. 3645–53.53.

- 62) Boesen et al study. The vascular response to activity
- 63) Boesen et al: Electro-coagulation of the Achilles
- 64) (Hydrodilatation (distension arthrography): a long-term clinical outcome series. Watson L, et al. Br J Sports Med. 2007 Mar;41(3):167-73. Epub 2006 Dec 18.).
- 65) Hydrodilatation in the management of shoulder capsulitis. Bell S, Coghlan J, Richardson M. Australas Radiol. 2003 Sep; 47(3):247-51.).
- 66) TREATMENT OF RIGID SHOULDERS BY JOINT DISTENSION DURING ARTHROGRAPHY.
ANDREN L, LUNDBERG BJ Acta Orthop Scand. 1965; 36:45-53.