

Referencer til artikel I Dansk Sportsmedicin nr. 3, 2014:

Patella tendinopati hos mandlig elite volleyballspiller - rehabilitering og tilbage til sport

Af fysioterapeut Lisbeth Lund Pedersen, Idrætsklinikken Sygehus Lillebælt, Middelfart

1. Magnusson S. Peter, Henning Langberg and Michael Kjaer. The pathogenesis of tendinopathy: balancing the response to loading. *Nat. Rev. Rheumatol.* 6, 262–268, doi:10.1038/nrrheum.2010.43, 2010.
2. Bloch, Thorlund Jonas. Patella tendinopati. *Fagforum For Idrætsfysioterapi.*
3. Cook J L, Purdam C R. Is tendon pathology a continuum? A pathology model to explain the clinical presentation of load-induced tendinopathy. *Br J Sports Med.* 43:409–416. doi:10.1136/bjism.2008.051193,, 2009.
4. Scott A, et al. Sports and exercise-related tendinopathies: a review of selected topical issues by participants of the second International Scientific Tendinopathy Symposium (ISTS) Vancouver 2012. *Br J Sports Med.* 47:536–544. doi:10.1136/bjsports-2013-092329, 2013.
5. Cook JL, et al. The challenge of managing tendinopathy in competing athletes. *Br J Sports Med.* 0:1–6. doi:10.1136/bjsports-2012-092078, 2013.
6. Fredberg U., K. Stengaard-Pedersen. Chronic tendinopathy tissue pathology, pain mechanisms, and etiology with a special focus on inflammation. *Scand J Med Sci Sports.* 18: 3–15, DOI: 10.1111/j.1600-0838.2007.00746.x, 2008.
7. Zwerver J, S W Bredeweg, A L Hof,. Biomechanical analysis of the single-leg decline squat. *Br J Sports Med.* 41:264–268. doi: 10.1136/bjism.2006.032482, 2007.
8. Bisseling RW, Hof AL, Bredeweg SW, et al. Relationship between landing strategy and patellar tendinopathy in volleyball. *Br J Sports Med.* 41:e8 (<http://www.bjsportmed.com/cgi/content/full/41/7/e8>). doi: 10.1136/bjism.2006.032565, 2007.
9. Malliaras P, Cook J. Reduced ankle dorsiflexion range may increase the risk of patellar tendon injury among volleyball players. *J Sci Med Sports.* 9(4):304 – 309, doi:10.1016/j.jsams.2006.03.015., 2000.
10. Reeser JC, Verhagen E, Briner WW, et al. Strategies for the prevention of volleyball related injuries. *r J Sports Med.* 40:594–600., 2006.
11. Thomee, R. A. Comprehensive Treatment Approach for Patellofemoral Pain Syndrome in Young Women. *PHYS THER.* 77:1690-1703, 1997.

12. Silbernagel, K, et al. Continued Sports Activity, Using a Pain-Monitoring Model, During Rehabilitation in Patients With Achilles Tendinopathy. A Randomized Controlled Study. *The American Journal of Sports Medicine*. Vol. 35, No. 6, DOI: 10.1177/0363546506298279, 2007.
13. Visnes, H., and R. Bahr. The evolution of eccentric training as treatment for patellar tendinopathy (jumper's knee): a critical review of exercise programmes. *Br. J. Sports Med*. 41:217-223, 2007.
14. Malliaras Peter, Christian J. Barton, Neil D. Reeves, Henning Langberg. Achilles and Patellar Tendinopathy Loading Programmes, A Systematic Review Comparing Clinical Outcomes and Identifying Potential Mechanisms for Effectiveness. *Sports Med*. 43:267–286, DOI 10.1007/s40279-013-0019-z, 2013.
15. Visnes, H, et al. No Effect of Eccentric Training on Jumper's Knee in Volleyball Players During the Competitive Season. A Randomized Clinical Trial. *Clin J Sport Med*. 15:227–234, 2005.
16. Langberg, H, et al. Eccentric rehabilitation exercise increases peritendinous type I collagen synthesis in humans with Achilles tendinosis. *Scand J Med Sci Sports*. 17: 61–66, DOI: 10.1111/j.1600-0838.2006.00522.x, 2007.
17. Kongsgaard M, Kovanen V, Aagaard P, et al. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scand J Med Sci Sports*. s. 19: 790-802, 2009.
18. Kongsgaard M, Klaus Qvortrup, Jytte Larsen, Per Aagaard, Simon Doessing, Philip Hansen, Michael Kjaer and S. Peter Magnusson. Fibril Morphology and Tendon Mechanical Properties in Patellar Tendinopathy: Effects of Heavy Slow Resistance Training. *Am J Sports Med*. 38: 749 originally published online February 12, 2010, DOI: 10.1177/0363546509350915, 2010.
19. Richards DP, Ajemian SV, Wiley JP, et al. Knee joint dynamics predict patellar tendinitis in elite volleyball players. *Am J Sports Med*. 24:676–683, 1996.
20. Edwards S, Steele JR, Cook JL, Purdam CR, McGhee DE, Munro BJ. Edwards S, Steele JR, Cook JL, Purdam CR, McGhee DE, Munro BJ. Characterizing patellar tendon loading during the landing phase of a stop-jump task. *Scand J Med Sci Sports*. 22: 2–11, doi: 10.1111/j.1600-0838.2010.01119.x,, 2012.
21. Visnes H, Bahr R. Training volume and body composition as risk factors for developing Jumper's knee among young elite volleyball players. *Scand J Med Sci Sports*. Published Online First: 20 January 2012. doi: 10.1111/j.1600-0838.2011.01430.x,, 2012.
22. Visnes H, et al. Jumper's knee paradox—jumping ability is a risk factor for developing jumper's knee: a 5-year prospective study. *Br J Sports Med*. 2013;47:503–507. doi:10.1136/bjsports-2012-091385,, 2013.
23. Visentini PJ, Khan KM, Cook JL, et al. The VISA score: an index of severity of symptoms in patients with jumper's knee (patellar tendinosis). *J Sci Med Sport*. 1:22–8, 1998.
24. Lian O, Engebretsen L, Bahr R. Prevalence of jumper's knee among elite athletes from different sports: a cross-sectional study. *Am J Sports Med*. Vol. 33, No. 4, DOI: 10.1177/0363546504270454, 2005.