

Referenceliste til artikel i Dansk Sportsmedicin nr. 3, 2007:

"Næring til muskeltvækst"

af cand.scient, PhD-studerende Mette Hansen

1. Andersen LL, Tufekovic G, Zebis MK, Cramer RM, Verlaan G, Kjaer M, Suetta C, Magnusson P and Aagaard P. The effect of resistance training combined with timed ingestion of protein on muscle fiber size and muscle strength. *Metabolism* 54: 151-156, 2005.
2. Anthony JC, Anthony TG, Kimball SR, Vary TC and Jefferson LS. Orally administered leucine stimulates protein synthesis in skeletal muscle of postabsorptive rats in association with increased eIF4F formation. *J Nutr* 130: 139-145, 2000.
3. Arnal MA, Mosoni L, Boirie Y, Houlier ML, Morin L, Verdier E, Ritz P, Antoine JM, Prugnaud J, Beaufriere B and Mirand PP. Protein pulse feeding improves protein retention in elderly women. *Am J Clin Nutr* 69: 1202-1208, 1999.
4. Arnal MA, Mosoni L, Boirie Y, Houlier ML, Morin L, Verdier E, Ritz P, Antoine JM, Prugnaud J, Beaufriere B and Mirand PP. Protein feeding pattern does not affect protein retention in young women. *J Nutr* 130: 1700-1704, 2000.
5. Biolo G, Maggi SP, Williams BD, Tipton KD and Wolfe RR. Increased rates of muscle protein turnover and amino acid transport after resistance exercise in humans. *Am J Physiol* 268: E514-E520, 1995.
6. Biolo G, Tipton KD, Klein S and Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol* 273: E122-E129, 1997.
7. Bohe J, Low JF, Wolfe RR and Rennie MJ. Latency and duration of stimulation of human muscle protein synthesis during continuous infusion of amino acids. *J Physiol* 532: 575-579, 2001.
8. Boirie Y, Dangin M, Gachon P, Vasson MP, Maubois JL and Beaufriere B. Slow and fast dietary proteins differently modulate postprandial protein accretion. *Proc Natl Acad Sci U S A* 94: 14930-14935, 1997.
9. Borsheim E, Cree MG, Tipton KD, Elliott TA, Aarsland A and Wolfe RR. Effect of carbohydrate intake on net muscle protein synthesis during recovery from resistance exercise. *J Appl Physiol* 96: 674-678, 2004.
10. Borsheim E, Tipton KD, Wolf SE and Wolfe RR. Essential amino acids and muscle protein recovery from resistance exercise. *Am J Physiol Endocrinol Metab* 283: E648-E657, 2002.
11. Bos C, Metges CC, Gaudichon C, Petzke KJ, Pueyo ME, Morens C, Everwand J, Benamouzig R and Tome D. Postprandial kinetics of dietary amino acids are the main determinant of their metabolism after soy or milk protein ingestion in humans. *J Nutr* 133: 1308-1315, 2003.
12. Bunout B, Barrera G, de la MP, Avendano M, Gattas V, Petermann M and Hirsh S. Effects of nutritional supplementation and resistance training on muscle strength in free living elders. Results of one year follow. *J Nutr Health Aging* 8: 68-75, 2004.
13. Campbell WW, Crim MC, Young VR, Joseph LJ and Evans WJ. Effects of resistance training and dietary protein intake on protein metabolism in older adults. *Am J Physiol* 268: E1143-E1153, 1995.
14. Candow DG, Chilibeck PD, Facci M, Abeysekara S and Zello GA. Protein supplementation before and after resistance training in older men. *Eur J Appl Physiol* 97: 548-556, 2006.

15. Carter JM, Bembien DA, Knehans AW, Bembien MG and Witten MS. Does nutritional supplementation influence adaptability of muscle to resistance training in men aged 48 to 72 years. *J Geriatr Phys Ther* 28: 40-47, 2005.
16. Cuthbertson D, Smith K, Babraj J, Leese G, Waddell T, Atherton P, Wackerhage H, Taylor PM and Rennie MJ. Anabolic signaling deficits underlie amino acid resistance of wasting, aging muscle. *FASEB J* 19: 422-424, 2005.
17. Dangin M, Boirie Y, Garcia-Rodenas C, Gachon P, Fauquant J, Callier P, Ballevre O and Beaufriere B. The digestion rate of protein is an independent regulating factor of postprandial protein retention. *Am J Physiol Endocrinol Metab* 280: E340-E348, 2001.
18. Dangin M, Guillet C, Garcia-Rodenas C, Gachon P, Bouteloup-Demange C, Reiffers-Magnani K, Fauquant J, Ballevre O and Beaufriere B. The rate of protein digestion affects protein gain differently during aging in humans. *J Physiol* 549: 635-644, 2003.
19. Dardevet D, Sornet C, Balage M and Grizard J. Stimulation of in vitro rat muscle protein synthesis by leucine decreases with age. *J Nutr* 130: 2630-2635, 2000.
20. Economos CD, Bortz SS and Nelson ME. Nutritional practices of elite athletes. Practical recommendations. *Sports Med* 16: 381-399, 1993.
21. Elliot TA, Cree MG, Sanford AP, Wolfe RR and Tipton KD. Milk ingestion stimulates net muscle protein synthesis following resistance exercise. *Med Sci Sports Exerc* 38: 667-674, 2006.
22. Esmarck B, Andersen JL, Olsen S, Richter EA, Mizuno M and Kjaer M. Timing of postexercise protein intake is important for muscle hypertrophy with resistance training in elderly humans. *J Physiol* 535: 301-311, 2001.
23. Godard MP, Williamson DL and Trappe SW. Oral amino-acid provision does not affect muscle strength or size gains in older men. *Med Sci Sports Exerc* 34: 1126-1131, 2002.
24. Guillet C, Prod'homme M, Balage M, Gachon P, Giraudet C, Morin L, Grizard J and Boirie Y. Impaired anabolic response of muscle protein synthesis is associated with S6K1 dysregulation in elderly humans. *FASEB J* 18: 1586-1587, 2004.
25. Halton TL and Hu FB. The effects of high protein diets on thermogenesis, satiety and weight loss: a critical review. *J Am Coll Nutr* 23: 373-385, 2004.
26. Hamalainen E, Adlercreutz H, Puska P and Pietinen P. Diet and serum sex hormones in healthy men. *J Steroid Biochem* 20: 459-464, 1984.
27. Hawley JA, Tipton KD and Millard-Stafford ML. Promoting training adaptations through nutritional interventions. *J Sports Sci* 24: 709-721, 2006.
28. Holm L, Esmarck B, Mizuno M, Hansen H, Suetta C, Holmich P, Krogsgaard M and Kjaer M. The effect of protein and carbohydrate supplementation on strength training outcome of rehabilitation in ACL patients. *J Orthop Res* .: 2006.
29. Hong SO and Layman DK. Effects of leucine on in vitro protein synthesis and degradation in rat skeletal muscles. *J Nutr* 114: 1204-1212, 1984.
30. Karlsson HK, Nilsson PA, Nilsson J, Chibalin AV, Zierath JR and Blomstrand E. Branched-chain amino acids increase p70S6k phosphorylation in human skeletal muscle after resistance exercise. *Am J Physiol Endocrinol Metab* 287: E1-E7, 2004.
31. Katsanos CS, Kobayashi H, Sheffield-Moore M, Aarsland A and Wolfe RR. Aging is associated with diminished accretion of muscle proteins after the ingestion of a small bolus of essential amino acids. *Am J Clin Nutr* 82: 1065-1073, 2005.
32. Katsanos CS, Kobayashi H, Sheffield-Moore M, Aarsland A and Wolfe RR. A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly. *Am J Physiol Endocrinol Metab* 2006.

33. Kerksick CM, Rasmussen CJ, Lancaster SL, Magu B, Smith P, Melton C, Greenwood M, Almada AL, Earnest CP and Kreider RB. The effects of protein and amino acid supplementation on performance and training adaptations during ten weeks of resistance training. *J Strength Cond Res* 20: 643-653, 2006.
34. Kimball SR. Regulation of global and specific mRNA translation by amino acids. *J Nutr* 132: 883-886, 2002.
35. Kimball SR, Shantz LM, Horetsky RL and Jefferson LS. Leucine regulates translation of specific mRNAs in L6 myoblasts through mTOR-mediated changes in availability of eIF4E and phosphorylation of ribosomal protein S6. *J Biol Chem* 274: 11647-11652, 1999.
36. Koopman R, Verdijk L, Manders RJ, Gijsen AP, Gorselink M, Pijpers E, Wagenmakers AJ and van Loon LJ. Co-ingestion of protein and leucine stimulates muscle protein synthesis rates to the same extent in young and elderly lean men. *Am J Clin Nutr* 84: 623-632, 2006.
37. Koopman R, Wagenmakers AJ, Manders RJ, Zorenc AH, Senden JM, Gorselink M, Keizer HA and van Loon LJ. Combined ingestion of protein and free leucine with carbohydrate increases postexercise muscle protein synthesis in vivo in male subjects. *Am J Physiol Endocrinol Metab* 288: E645-E653, 2005.
38. Lambert CP, Frank LL and Evans WJ. Macronutrient considerations for the sport of bodybuilding. *Sports Med* 34: 317-327, 2004.
39. Mikkelsen PB, Toubro S and Astrup A. Effect of fat-reduced diets on 24-h energy expenditure: comparisons between animal protein, vegetable protein, and carbohydrate. *Am J Clin Nutr* 72: 1135-1141, 2000.
40. Miller SL, Tipton KD, Chinkes DL, Wolf SE and Wolfe RR. Independent and combined effects of amino acids and glucose after resistance exercise. *Med Sci Sports Exerc* 35: 449-455, 2003.
41. Moore DR, Del Bel NC, Nizi KI, Hartman JW, Tang JE, Armstrong D and Phillips SM. Resistance training reduces fasted- and fed-state leucine turnover and increases dietary nitrogen retention in previously untrained young men. *J Nutr* 137: 985-991, 2007.
42. Morens C, Bos C, Pueyo ME, Benamouzig R, Gausseres N, Luengo C, Tome D and Gaudichon C. Increasing habitual protein intake accentuates differences in postprandial dietary nitrogen utilization between protein sources in humans. *J Nutr* 133: 2733-2740, 2003.
43. Mourao DM, Bressan J, Campbell WW and Mattes RD. Effects of food form on appetite and energy intake in lean and obese young adults. *Int J Obes (Lond)* 2007.
44. Nordic Council of Ministers and The Nordic Food Policy Co-operation. *Nordic Nutrition Recommendations 2004 (Nord 2004:013)*. 2004.
45. Ref Type: Internet Communication
46. Phillips SM, Hartman JW and Wilkinson SB. Dietary protein to support anabolism with resistance exercise in young men. *J Am Coll Nutr* 24: 134S-139S, 2005.
47. Phillips SM, Parise G, Roy BD, Tipton KD, Wolfe RR and Tamopolsky MA. Resistance-training-induced adaptations in skeletal muscle protein turnover in the fed state. *Can J Physiol Pharmacol* 80: 1045-1053, 2002.
48. Phillips SM, Tipton KD, Aarsland A, Wolf SE and Wolfe RR. Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *Am J Physiol* 273: E99-107, 1997.
49. Phillips SM, Tipton KD, Ferrando AA and Wolfe RR. Resistance training reduces the acute exercise-induced increase in muscle protein turnover. *Am J Physiol* 276: E118-E124, 1999.
50. Rasmussen BB, Tipton KD, Miller SL, Wolf SE and Wolfe RR. An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J Appl Physiol* 88: 386-392, 2000.
51. Rennie MJ, Bohe J and Wolfe RR. Latency, duration and dose response relationships of amino acid effects on human muscle protein synthesis. *J Nutr* 132: 3225S-3227S, 2002.

52. Rieu I, Balage M, Sornet C, Giraudet C, Pujos E, Grizard J, Mosoni L and Dardevet D. Leucine supplementation improves muscle protein synthesis in elderly men independently of hyperaminoacidaemia. *J Physiol* 575: 305-315, 2006.
53. Rosendahl E, Lindelof N, Littbrand H, Yifter-Lindgren E, Lundin-Olsson L, Haglin L, Gustafson Y and Nyberg L. High-intensity functional exercise program and protein-enriched energy supplement for older persons dependent in activities of daily living: a randomised controlled trial. *Aust J Physiother* 52: 105-113, 2006.
54. Roy BD, Tarnopolsky MA, MacDougall JD, Fowles J and Yarasheski KE. Effect of glucose supplement timing on protein metabolism after resistance training. *J Appl Physiol* 82: 1882-1888, 1997.
55. Tarnopolsky MA, Atkinson SA, MacDougall JD, Chesley A, Phillips S and Schwarcz HP. Evaluation of protein requirements for trained strength athletes. *J Appl Physiol* 73: 1986-1995, 1992.
56. Tipton KD, Borsheim E, Wolf SE, Sanford AP and Wolfe RR. Acute response of net muscle protein balance reflects 24-h balance after exercise and amino acid ingestion. *Am J Physiol Endocrinol Metab* 284: E76-E89, 2003.
57. Tipton KD, Elliott TA, Cree MG, Aarsland AA, Sanford AP and Wolfe RR. Stimulation of net muscle protein synthesis by whey protein ingestion before and after exercise. *Am J Physiol Endocrinol Metab* 292: E71-E76, 2007.
58. Tipton KD, Elliott TA, Cree MG, Wolf SE, Sanford AP and Wolfe RR. Ingestion of casein and whey proteins result in muscle anabolism after resistance exercise. *Med Sci Sports Exerc* 36: 2073-2081, 2004.
59. Tipton KD, Ferrando AA, Phillips SM, Doyle D, Jr. and Wolfe RR. Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am J Physiol* 276: E628-E634, 1999.
60. Tipton KD, Rasmussen BB, Miller SL, Wolf SE, Owens-Stovall SK, Petrini BE and Wolfe RR. Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrinol Metab* 281: E197-E206, 2001.
61. Tipton KD and Wolfe RR. Protein and amino acids for athletes. *J Sports Sci* 22: 65-79, 2004.
62. Todd KS, Butterfield GE and Calloway DH. Nitrogen balance in men with adequate and deficient energy intake at three levels of work. *J Nutr* 114: 2107-2118, 1984.
63. Tome D and Bos C. Dietary protein and nitrogen utilization. *J Nutr* 130: 1868S-1873S, 2000.
64. Volpi E, Mittendorfer B, Rasmussen BB and Wolfe RR. The response of muscle protein anabolism to combined hyperaminoacidemia and glucose-induced hyperinsulinemia is impaired in the elderly. *J Clin Endocrinol Metab* 85: 4481-4490, 2000.
65. Welle S, Thornton C, Statt M and McHenry B. Postprandial myofibrillar and whole body protein synthesis in young and old human subjects. *Am J Physiol* 267: E599-E604, 1994.
66. Welle S and Thornton CA. High-protein meals do not enhance myofibrillar synthesis after resistance exercise in 62- to 75-yr-old men and women. *Am J Physiol* 274: E677-E683, 1998.
67. Wilkinson SB, Tarnopolsky MA, Macdonald MJ, Macdonald JR, Armstrong D and Phillips SM. Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *Am J Clin Nutr* 85: 1031-1040, 2007.
68. Wolfe RR. Skeletal muscle protein metabolism and resistance exercise. *J Nutr* 136: 525S-528S, 2006.