

## **Referencer til artikel i Dansk Sportsmedicin nr. 2, 2010:**

### **"Risikofaktorer for udvikling af patellofemoralt smertesyndrom og medialt tibialt stress-syndrom"**

*Af Michael skovdal Rathleff og Camilla R. Knudsen*

- (1) Witvrouw E, Werner S, Mikkelsen C, Van TD, Vanden BL, Cerulli G. Clinical classification of patellofemoral pain syndrome: guidelines for non-operative treatment. *Knee Surg Sports Traumatol Arthrosc* 2005 Mar;13(2):122-30.
- (2) Cowan SM, Bennell KL, Hodges PW, Crossley KM, McConnell J. Delayed onset of electromyographic activity of vastus medialis obliquus relative to vastus lateralis in subjects with patellofemoral pain syndrome. *Arch Phys Med Rehabil* 2001 Feb;82(2):183-9.
- (3) Kortebéin PM, Kaufman KR, Basford JR, Stuart MJ. Medial tibial stress syndrome. *Med Sci Sports Exerc* 2000 Mar;32(3 Suppl):S27-S33.
- (4) Van TD, Cowan S, Coorevits P, Duvigneaud N, Witvrouw E. Delayed vastus medialis obliquus to vastus lateralis onset timing contributes to the development of patellofemoral pain in previously healthy men: a prospective study. *Am J Sports Med* 2009 Jun;37(6):1099-105.
- (5) Duvigneaud N, Barnard E, Stevens V, Witvrouw E, Van TD. Isokinetic assessment of patellofemoral pain syndrome: A prospective study in female recruits. *Isokinetics and Exercise Science* 2008;16:213-9.
- (6) Boling MC, Padua DA, Marshall SW, Guskiewicz K, Pyne S, Beutler A. A prospective investigation of biomechanical risk factors for patellofemoral pain syndrome: the Joint Undertaking to Monitor and Prevent ACL Injury (JUMP-ACL) cohort. *Am J Sports Med* 2009 Nov;37(11):2108-16.
- (7) Thijs Y, De CD, Roosen P, Witvrouw E. Gait-related intrinsic risk factors for patellofemoral pain in novice recreational runners. *Br J Sports Med* 2008 Jun;42(6):466-71.
- (8) Witvrouw E, Lysens R, Bellemans J, Cambier D, Vanderstraeten G. Intrinsic risk factors for the development of anterior knee pain in an athletic population. A two-year prospective study. *Am J Sports Med* 2000 Jul;28(4):480-9.
- (9) Clement DB, Taunton JE, Smart GW, McNicol KL. A survey of overuse running injuries. *Phys Sports Med* 1981;9:47-58.
- (10) Yates B, White S. The incidence and risk factors in the development of medial tibial stress syndrome among naval recruits. *Am J Sports Med* 2004 Apr;32(3):772-80.
- (11) Willems TM, De CD, Delbaere K, Vanderstraeten G, De CA, Witvrouw E. A prospective study of gait related risk factors for exercise-related lower leg pain. *Gait Posture* 2006 Jan;23(1):91-8.
- (12) Willems TM, Witvrouw E, De CA, De CD. Gait-related risk factors for exercise-related lower-leg pain during shod running. *Med Sci Sports Exerc* 2007 Feb;39(2):330-9.

- (13) Reinking MF. Exercise-related leg pain in female collegiate athletes: the influence of intrinsic and extrinsic factors. *Am J Sports Med* 2006 Sep;34(9):1500-7.
- (14) Plisky MS, Rauh MJ, Heiderscheit B, Underwood FB, Tank RT. Medial tibial stress syndrome in high school cross-country runners: incidence and risk factors. *J Orthop Sports Phys Ther* 2007 Feb;37(2):40-7.
- (15) Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther* 2010 Feb;40(2):42-51.
- (16) Feller JA, Amis AA, Andrich JT, Arendt EA, Erasmus PJ, Powers CM. Surgical biomechanics of the patellofemoral joint. *Arthroscopy* 2007 May;23(5):542-53.
- (17) Cowan SM, Bennell KL, Crossley KM, Hodges PW, McConnell J. Physical therapy alters recruitment of the vasti in patellofemoral pain syndrome. *Med Sci Sports Exerc* 2002 Dec;34(12):1879-85.
- (18) Bouche RT, Johnson CH. Medial tibial stress syndrome (tibial fasciitis): a proposed pathomechanical model involving fascial traction. *J Am Podiatr Med Assoc* 2007 Jan;97(1):31-6.
- (19) Mills K, Blanch P, Chapman AR, McPoil TG, Vicenzino B. Foot Orthoses and Gait: A Systematic Review and Meta-analysis of Literature Pertaining to Potential Mechanisms. *Br J Sports Med* 2009 Dec 8.